

Adsorption of glyphosate to Cambisols, Podzol and silica sand

Axel Mentler¹, M. Paredes^{1,2}, M. Fuerhacker²

University of Natural Resources and Applied Life Sciences, Vienna BOKU,

¹ *Institute of Soil Science* ² *Institute of Sanitary Engineering and Water Pollution Control*
Ass.Prof. DI Dr. Axel MENTLER, University of Natural Resources and Applied Life Sciences
Vienna, Department of Forest- and Soil Sciences, Peter Jordanstr. 82, A-1190 Vienna,
Austria e-mail: axel.mentler@boku.ac.at

The glyphosate is one of the most applied organic herbicides with increasing importance nowadays. Glyphosate is a post-emergency non-selective broad spectrum herbicide extensively used in agriculture for the control of most annual and perennial plants. It controls weeds by inhibiting the synthesis of aromatic amino acids necessary for protein formation in susceptible plants. The sorption of glyphosate by soils occurs due to the inner sphere complex formation with metals of soil oxides, which are related to the soil phosphate adsorption capacity. Several binding mechanisms have been suggested for the sorption of this herbicide, such as: electrostatic bonds in extremely acid, hydrogen bonds with humic substances and, especially, covalent bonds with Fe and Al oxides.

The present study should contribute to answer the question if glyphosate binds to different soil types and how various iron oxides contribute to the binding mechanisms.

Batch adsorption experiments were conducted according to Part C 18 (EU, 2001) with three different soils and silica sand for a comparison:

- Soil originating from Wienerwald (Orthic Podzol, Bs, FAO classification 1994)
- Eurosoil 7 (E7) (Dystric Cambisol, Ah, FAO classification 1994)
- Soil originating from Phyra (Dystric Cambisol, Ah, FAO classification 1994)
- Silica sand (acid washed)

The substances were measured either as phosphorous (P) after filtration by ICP-OES or/and as single compounds with HPLC/FLD after post column derivatisation with fluorescence detection (Pickering system).

The adsorption capacities of the two cambisols (Eurosoil (KD 188 – 299 l/kg) and Phyra (KD 13.8 – 29.3 l/kg)) were quite different and varied about one order of magnitude, although their Fe content was comparable. The Podzol (KD 467 – 519 l/kg) contained about 3 % soluble iron oxides and produced best adsorption results. The KD of silica sand was negligible. In the Wienerwald sample (Podzol) the main adsorption process seems to be based on iron oxides as the organic carbon content is quite low. Our data are in the range of those found in the literature; nevertheless the variations of the KD- values in different soils are significant and not only dependent of the Fe-content. A reduced sorption capacity could result in a leaching of glyphosate to groundwater.

Glyphosate can be adsorbed by the different substrates studied in different ways (e.g. Cambisols). Therefore, the use of glyphosate should be adapted to physico-chemical soil properties. The results in the literature show that iron oxides play an important role in the retention of glyphosate. In our study (data not shown) we could observe that from the investigated iron oxides (ferrihydrite, hematite and goethite) ferrihydrite had the highest impact on the adsorption process.